

# The Bohr–Pál Theorem and the Sobolev Space $W_2^{1/2}$

Vladimir Lebedev

**Abstract.** The well-known Bohr–Pál theorem asserts that for every continuous real-valued function  $f$  on the circle  $\mathbb{T}$  there exists a change of variable, i.e., a homeomorphism  $h$  of  $\mathbb{T}$  onto itself, such that the Fourier series of the superposition  $f \circ h$  converges uniformly. Subsequent improvements of this result imply that actually there exists a homeomorphism that brings  $f$  into the Sobolev space  $W_2^{1/2}(\mathbb{T})$ . This refined version of the Bohr–Pál theorem does not extend to complex-valued functions. We show that if  $\alpha < 1/2$ , then there exists a complex-valued  $f$  that satisfies the Lipschitz condition of order  $\alpha$  and at the same time has the property that  $f \circ h \notin W_2^{1/2}(\mathbb{T})$  for every homeomorphism  $h$  of  $\mathbb{T}$ .

2010 *Mathematics Subject Classification*: 42A16.

*Key words and phrases*: harmonic analysis, homeomorphisms of the circle, superposition operators, Sobolev spaces.

## 1. Introduction

For an arbitrary integrable function  $f$  on the circle  $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$  (where  $\mathbb{R}$  is the real line and  $\mathbb{Z}$  is the group of integers) consider its Fourier series:

$$f(t) \sim \sum_{k \in \mathbb{Z}} \widehat{f}(k) e^{ikt}, \quad t \in \mathbb{T}.$$

Recall that the Sobolev space  $W_2^{1/2}(\mathbb{T})$  is the space of all (integrable) functions  $f$  with

$$\sum_{k \in \mathbb{Z}} |\widehat{f}(k)|^2 |k| < \infty.$$

Let  $C(\mathbb{T})$  be the space of all continuous functions on  $\mathbb{T}$ .

The well-known Bohr–Pál theorem states that for every real-valued function  $f \in C(\mathbb{T})$  there exists a homeomorphism  $h$  of the circle  $\mathbb{T}$  onto itself, such that the superposition  $f \circ h$  belongs to the space  $U(\mathbb{T})$  of uniformly convergent Fourier series. (The theorem was obtained in a somewhat weaker form by J. Pál in [11], and in the final form by H. Bohr in [2].) The original

method of proof of this result uses conformal mappings and in fact allows (see [9, Sec. 3]) to obtain the following representation:

$$f \circ h = g + \psi, \quad g \in W_2^{1/2} \cap C(\mathbb{T}), \quad \psi \in V \cap C(\mathbb{T}), \quad (1)$$

where  $V(\mathbb{T})$  is the space of functions of bounded variation on  $\mathbb{T}$ . It is well-known that both  $W_2^{1/2} \cap C(\mathbb{T})$  and  $V \cap C(\mathbb{T})$  are subsets of  $U(\mathbb{T})$ , thus (1) implies  $f \circ h \in U(\mathbb{T})$ .

A substantial improvement of the Bohr–Pál theorem was obtained by A. A. Sahakian [12, Corollary 1], who showed that if  $a(n), n = 0, 1, 2, \dots$ , is a given positive sequence satisfying the condition  $\sum_n a(n) = \infty$  and a certain condition of regularity, then for every real-valued  $f \in C(\mathbb{T})$  there is a homeomorphism  $h$  such that  $\widehat{f \circ h}(k) = O(a(|k|))$ . An immediate consequence of Sahakian's result is that the term  $\psi$  in (1) can be omitted, i.e., the following refined version of the Bohr–Pál theorem holds: for every real-valued  $f \in C(\mathbb{T})$  there exists a homeomorphism  $h$  of  $\mathbb{T}$  onto itself, such that  $f \circ h \in W_2^{1/2}(\mathbb{T})$ . This refined version also follows from a result on conjugate functions, obtained by W. Jurkat and D. Waterman in [4] (see also [3, Theorem 9.5]). We note that Sahakian's result is obtained by purely real analysis technique whereas Jurkat and Waterman use an approach similar to the one used by Bohr and Pál. A very short proof of the refined version of the Bohr–Pál theorem was communicated to the author by A. Olevskiĭ, see [7, Sec. 3].

Another improvement of the Bohr–Pál theorem was obtained by J.-P. Kahane and Y. Katznelson [6] (see also [9], [5]). These authors showed that if  $K$  is a compact family of functions in  $C(\mathbb{T})$ , then there exists a homeomorphism  $h$  of  $\mathbb{T}$  such that  $f \circ h \in U(\mathbb{T})$  for all  $f \in K$ . This result naturally leads to a question if it is possible to attain the condition  $f \circ h \in W_2^{1/2}(\mathbb{T})$  for all  $f \in K$ . This question was posed by A. Olevskiĭ in [10]. A negative answer was obtained by the author of this work in [7, Theorem 4], it turns out that, given a real-valued  $u \in C(\mathbb{T})$ , the property that for every real-valued  $v \in C(\mathbb{T})$  there is a homeomorphism  $h$  such that both  $u \circ h$  and  $v \circ h$  are in  $W_2^{1/2}(\mathbb{T})$  is equivalent to the boundness of variation of  $u$ . Thus, in general, there is no single change of variable which will bring two real-valued functions in  $C(\mathbb{T})$  into  $W_2^{1/2}(\mathbb{T})$ . Certainly this amounts to the existence of a complex-valued  $f \in C(\mathbb{T})$  such that  $f \circ h \notin W_2^{1/2}(\mathbb{T})$  for every homeomorphism  $h$  of  $\mathbb{T}$ .

The purpose of this work is to show that there exists a complex-valued function  $f$  that is *very smooth* but at the same time has the property that  $f \circ h \notin W_2^{1/2}(\mathbb{T})$  for every homeomorphism  $h$  of  $\mathbb{T}$ .

Note that, as one can easily verify (see, e.g., [7], Sec. 3), the following two semi-norms

$$\begin{aligned} \|f\|_{W_2^{1/2}(\mathbb{T})} &= \left( \sum_{k \in \mathbb{Z}} |\widehat{f}(k)|^2 |k| \right)^{1/2}, \\ |||f|||_{W_2^{1/2}(\mathbb{T})} &= \left( \int_0^{2\pi} \frac{1}{\theta^2} \int_0^{2\pi} |f(t+\theta) - f(t)|^2 dt d\theta \right)^{1/2} \end{aligned} \quad (2)$$

are equivalent semi-norms on  $W_2^{1/2}(\mathbb{T})$ , i.e.,  $f$  is in  $W_2^{1/2}(\mathbb{T})$  if and only if  $|||f|||_{W_2^{1/2}(\mathbb{T})} < \infty$ , and  $c_1 \|f\|_{W_2^{1/2}(\mathbb{T})} \leq |||f|||_{W_2^{1/2}(\mathbb{T})} \leq c_2 \|f\|_{W_2^{1/2}(\mathbb{T})}$  for all  $f \in W_2^{1/2}(\mathbb{T})$ , where  $c_1, c_2 > 0$  do not depend on  $f$ . Thus, we see that every function that satisfies the Lipschitz condition of order greater than  $1/2$  belongs to  $W_2^{1/2}(\mathbb{T})$ . We shall show that, in general, there is no change of variable which will bring a complex-valued function that satisfies the Lipschitz condition of order less than  $1/2$  into  $W_2^{1/2}(\mathbb{T})$ . The author does not know if the same holds for the functions satisfying the Lipschitz condition of order  $1/2$  (see Remarks at the end of the paper).

## 2. Result

Let  $\omega$  be a modulus of continuity, i.e., a nondecreasing continuous function on  $[0, +\infty)$  such that  $\omega(0) = 0$  and  $\omega(x+y) \leq \omega(x) + \omega(y)$ . By  $\text{Lip}_\omega(\mathbb{T})$  we denote the class of all complex-valued functions  $f$  on  $\mathbb{T}$  with  $\omega(f, \delta) = O(\omega(\delta))$ ,  $\delta \rightarrow +0$ , where

$$\omega(f, \delta) = \sup_{|t_1 - t_2| \leq \delta} |f(t_1) - f(t_2)|, \quad \delta \geq 0,$$

is the modulus of continuity of  $f$ . For  $0 < \alpha \leq 1$  we just write  $\text{Lip}_\alpha$  instead of  $\text{Lip}_{\delta^\alpha}$ .

**Theorem.** *Suppose that  $\limsup_{\delta \rightarrow +0} \omega(\delta)/\sqrt{\delta} = \infty$ . Then there exists a complex-valued function  $f \in \text{Lip}_\omega(\mathbb{T})$  such that  $f \circ h \notin W_2^{1/2}(\mathbb{T})$  for every homeomorphism  $h$  of the circle  $\mathbb{T}$  onto itself. In particular, if  $\alpha < 1/2$ , then there exists a function of class  $\text{Lip}_\alpha(\mathbb{T})$  with this property.*

Ideologically the method of the proof of this theorem is close to the one used by the author to prove Theorem 4 in [7].

We shall need certain preliminary constructions and lemmas. Simple Lemma 1 below is purely technical.

**Lemma 1.** *Under the assumption of the theorem on  $\omega$  there exists a sequence  $\delta_k > 0$ ,  $k = 1, 2, \dots$ , such that*

$$\sum_{k=1}^{\infty} \delta_k < 2\pi/6, \quad (3)$$

$$\sum_{k=1}^{\infty} (\omega(\delta_k))^2 = \infty. \quad (4)$$

*Proof.* For each  $j = 1, 2, \dots$  we can find  $\varepsilon_j$  so that  $0 < \varepsilon_j < 2^{-(j+1)}$  and

$$\frac{(\omega(\varepsilon_j))^2}{\varepsilon_j} \geq 2^j.$$

Chose positive integers  $n_j$  satisfying

$$\frac{1}{2^{j+1}\varepsilon_j} \leq n_j < \frac{1}{2^j\varepsilon_j}, \quad j = 1, 2, \dots$$

Let  $N_0 = 1$  and let  $N_j = N_{j-1} + n_j$  for  $j = 1, 2, \dots$ . We define the sequence  $\delta_k$ ,  $k = 1, 2, \dots$ , by setting  $\delta_k = \varepsilon_j$  if  $N_{j-1} \leq k < N_j$ ,  $j = 1, 2, \dots$ . This yields

$$\sum_{k=1}^{\infty} \delta_k = \sum_{j=1}^{\infty} \sum_{N_{j-1} \leq k < N_j} \delta_k = \sum_{j=1}^{\infty} n_j \varepsilon_j \leq \sum_{j=1}^{\infty} \frac{1}{2^j} = 1,$$

and at the same time

$$\sum_{N_{j-1} \leq k < N_j} (\omega(\delta_k))^2 = n_j (\omega(\varepsilon_j))^2 \geq n_j \varepsilon_j 2^j \geq \frac{1}{2}.$$

The lemma is proved.

For a closed interval  $I = [a, b] \subseteq (0, 2\pi)$  let  $\Delta_I$  denote the “triangle” function supported on  $I$ , i.e., a continuous function on the interval  $[0, 2\pi]$  such that  $\Delta_I(t) = 0$  for all  $t \in [0, a] \cup [b, 2\pi]$ ,  $\Delta_I(c) = 1$ , where  $c = (a + b)/2$  is the center of  $I$ , and  $\Delta_I$  is linear on  $[a, c]$  and on  $[c, b]$ .

Let  $\delta_k, k = 1, 2, \dots$ , be the sequence from Lemma 1. Consider intervals  $I_k = [a_k, b_k] \subseteq (0, 2\pi)$  of length  $b_k - a_k = 6\delta_k$ , where  $a_k < b_k < a_{k+1}$ ,  $k = 1, 2, \dots$  (see (3)). For each  $k$  let  $J_k$  denote the left half of  $I_k$ , i.e.,  $J_k = [a_k, (a_k + b_k)/2]$ ,  $k = 1, 2, \dots$ .

Everywhere below we use  $u$  and  $v$  to denote two real-valued functions on  $\mathbb{T}$  defined by

$$u(t) = \sum_{k=1}^{\infty} \omega(\delta_k) \Delta_{I_k}(t), \quad v(t) = \sum_{k=1}^{\infty} \omega(\delta_k) \Delta_{J_k}(t), \quad t \in [0, 2\pi].$$

We shall show that the function  $f = u + iv$  satisfies the assertion of the theorem.

**Lemma 2.** *The functions  $u$  and  $v$  are of class  $\text{Lip}_\omega(\mathbb{T})$ .*

*Proof.* It is clear that for an arbitrary (closed) interval  $I \subseteq (0, 2\pi)$ , the function  $\Delta_I$  satisfies

$$|\Delta_I(t_1) - \Delta_I(t_2)| \leq \frac{2}{|I|} |t_1 - t_2|, \quad \text{for all } t_1, t_2 \in [0, 2\pi], \quad (5)$$

where  $|I|$  is the length of  $I$ .

Note also that if  $0 < x \leq y$ , then  $\omega(y)/y \leq 2\omega(x)/x$ . Indeed, let  $n = [y/x] + 1$ , where  $[\alpha]$  denotes the integer part of a number  $\alpha$ , then we have  $y \leq nx \leq 2y$ , so

$$\frac{\omega(y)}{y} \leq \frac{\omega(nx)}{y} \leq \frac{n\omega(x)}{y} \leq 2\frac{\omega(x)}{x}.$$

Let us show that  $u \in \text{Lip}_\omega(\mathbb{T})$ ; for  $v$  the proof is similar. It is easy to see that to prove the inclusion  $u \in \text{Lip}_\omega(\mathbb{T})$  it suffices to verify that for all  $t_1, t_2 \in \bigcup_k I_k$  we have

$$|u(t_1) - u(t_2)| \leq c\omega(|t_1 - t_2|),$$

where  $c > 0$  does not depend on  $t_1$  and  $t_2$ .

First we consider the case when  $t_1$  and  $t_2$  belong to the same interval  $I_k$ . If that is the case, then, since  $|t_1 - t_2| \leq |I_k| = 6\delta_k$ , we have

$$\frac{\omega(6\delta_k)}{6\delta_k} \leq 2 \frac{\omega(|t_1 - t_2|)}{|t_1 - t_2|},$$

so (see (5)),

$$\begin{aligned} |u(t_1) - u(t_2)| &= \omega(\delta_k) |\Delta_{I_k}(t_1) - \Delta_{I_k}(t_2)| \leq \\ &\leq \omega(\delta_k) \frac{2}{6\delta_k} |t_1 - t_2| \leq 2 \frac{\omega(6\delta_k)}{6\delta_k} |t_1 - t_2| \leq 4\omega(|t_1 - t_2|). \end{aligned}$$

Consider now the case when  $t_1 \in I_{k_1}$ ,  $t_2 \in I_{k_2}$ ,  $k_1 \neq k_2$ . We can assume that  $t_1 < t_2$ , and hence  $0 < t_1 < b_{k_1} < a_{k_2} < t_2 < 2\pi$ . Using the previous estimate, we obtain

$$|u(t_1) - u(t_2)| \leq |u(t_1)| + |u(t_2)| = |u(t_1) - u(b_{k_1})| + |u(t_2) - u(a_{k_2})| \leq 8\omega(|t_1 - t_2|).$$

The lemma is proved.

For  $n = 1, 2, \dots$  we define functions  $u_n$  by

$$u_n(t) = \max\{u(t), 1/n\}, \quad t \in \mathbb{T}.$$

As above,  $V(\mathbb{T})$  stands for the class of functions of bounded variation on  $\mathbb{T}$ .

**Lemma 3.** *The functions  $u_n$ ,  $n = 1, 2, \dots$ , have the following properties:*

$$|u_n(t_1) - u_n(t_2)| \leq |u(t_1) - u(t_2)| \quad \text{for all } t_1, t_2 \in \mathbb{T} \quad \text{and all } n; \quad (6)$$

$$u_n \in V(\mathbb{T}) \quad \text{for all } n; \quad (7)$$

$$\sup_n \left| \int_{\mathbb{T}} v(t) du_n(t) \right| = \infty. \quad (8)$$

*Proof.* Properties (6) and (7) are obvious. Let us verify (8). To this end consider the middle thirds of the intervals  $J_k$ , namely, the intervals  $J_k^* = [a_k + \delta_k, a_k + 2\delta_k]$ ,  $k = 1, 2, \dots$ . Note that if

$$\frac{\omega(\delta_k)}{3} \geq \frac{1}{n}, \quad (9)$$

then the function  $u_n$  coincides with  $u$  on  $J_k^*$ . So, if (9) holds, then  $u_n$  is monotonically increasing on  $J_k^*$ , and for its values at the endpoints of  $J_k^*$  we have

$$u_n(a_k + \delta_k) = \omega(\delta_k)/3, \quad u_n(a_k + 2\delta_k) = 2\omega(\delta_k)/3.$$

It is easily seen, that for each  $k$

$$\min_{J_k^*} v = 2\omega(\delta_k)/3.$$

Taking into account that  $u$ , and hence  $u_n$ , is non-decreasing on each interval  $J_k$ , we see that for all  $n$  and  $k$  satisfying condition (9)

$$\int_{J_k} v du_n \geq \int_{a_k + \delta_k}^{a_k + 2\delta_k} v du_n \geq \frac{2}{3}\omega(\delta_k) \int_{a_k + \delta_k}^{a_k + 2\delta_k} du_n = \frac{2}{3}\omega(\delta_k) \frac{1}{3}\omega(\delta_k) = \frac{2}{9}(\omega(\delta_k))^2.$$

In addition (since  $u_n$  is non-decreasing on each  $J_k$ ) we have

$$\int_{J_k} v du_n \geq 0$$

for all  $n$  and  $k$ . Thus, taking into account that  $v$  vanishes outside  $\bigcup_{k=1}^{\infty} J_k$ , we obtain

$$\int_{\mathbb{T}} v du_n = \sum_{k=1}^{\infty} \int_{J_k} v du_n \geq \sum_{k: \omega(\delta_k) \geq 3/n} \int_{J_k} v du_n \geq \sum_{k: \omega(\delta_k) \geq 3/n} \frac{2}{9}(\omega(\delta_k))^2.$$

Applying (4) we see that (8) holds. The lemma is proved.

We shall also need the following auxiliary lemma.

**Lemma 4.** *If  $x, y \in W_2^{1/2} \cap C(\mathbb{T})$  and  $y \in V(\mathbb{T})$ , then*

$$\left| \frac{1}{2\pi} \int_{\mathbb{T}} x(t) dy(t) \right| \leq \|x\|_{W_2^{1/2}(\mathbb{T})} \|y\|_{W_2^{1/2}(\mathbb{T})}.$$

*Proof.* Integration by parts yields

$$\frac{1}{2\pi} \int_0^{2\pi} e^{ikt} dy(t) = -\frac{1}{2\pi} \int_0^{2\pi} y(t) de^{ikt} = -ik\widehat{y}(-k).$$

So, if  $x$  is a trigonometric polynomial, then, using Cauchy inequality, we obtain

$$\begin{aligned} \left| \frac{1}{2\pi} \int_{\mathbb{T}} x(t) dy(t) \right| &= \left| \sum_k \widehat{x}(k) \frac{1}{2\pi} \int_{\mathbb{T}} e^{ikt} dy(t) \right| = \\ &= \left| \sum_k \widehat{x}(k) (-ik) \widehat{y}(-k) \right| \leq \|x\|_{W_2^{1/2}(\mathbb{T})} \|y\|_{W_2^{1/2}(\mathbb{T})}. \end{aligned}$$

To see that the assertion of the lemma holds in the general case, consider the Fejér sums  $\sigma_N(x)$  of the function  $x$ :

$$\sigma_N(x)(t) = \sum_{|k| \leq N} \left(1 - \frac{|k|}{N}\right) \widehat{x}(k) e^{ikt}.$$

Since  $|\widehat{\sigma_N(x)}(k)| \leq |\widehat{x}(k)|$  for all  $k \in \mathbb{Z}$ , we have  $\|\sigma_N(x)\|_{W_2^{1/2}(\mathbb{T})} \leq \|x\|_{W_2^{1/2}(\mathbb{T})}$ . Hence,

$$\left| \frac{1}{2\pi} \int_{\mathbb{T}} \sigma_N(x)(t) dy(t) \right| \leq \|\sigma_N(x)\|_{W_2^{1/2}(\mathbb{T})} \|y\|_{W_2^{1/2}(\mathbb{T})} \leq \|x\|_{W_2^{1/2}(\mathbb{T})} \|y\|_{W_2^{1/2}(\mathbb{T})}.$$

At the same time, since  $y$  is of bounded variation and  $\sigma_N(x)$  converges uniformly to  $x$  it is clear that

$$\frac{1}{2\pi} \int_{\mathbb{T}} \sigma_N(x)(t) dy(t) \rightarrow \frac{1}{2\pi} \int_{\mathbb{T}} x(t) dy(t)$$

as  $N \rightarrow \infty$ . The lemma is proved.

Now we proceed directly to the proof of the theorem. Let  $f = u + iv$ . Lemma 2 yields  $f \in \text{Lip}_\omega(\mathbb{T})$ , so it remains to show that  $f \circ h \notin W_2^{1/2}(\mathbb{T})$  for every homeomorphism  $h$  of  $\mathbb{T}$ . It is obvious that if a function is in  $W_2^{1/2}(\mathbb{T})$ , then both its real and imaginary parts are in  $W_2^{1/2}(\mathbb{T})$  as well. Assume that, contrary to the assertion of the theorem,  $f \circ h \in W_2^{1/2}(\mathbb{T})$  for a certain homeomorphism  $h$ . Then we have  $u \circ h \in W_2^{1/2}(\mathbb{T})$  and  $v \circ h \in W_2^{1/2}(\mathbb{T})$ .

Note that (6) implies  $|u_n \circ h(t_1) - u_n \circ h(t_2)| \leq |u \circ h(t_1) - u \circ h(t_2)|$  for all  $t_1, t_2 \in \mathbb{T}$ . Using the equivalence of the semi-norms  $\|\cdot\|_{W_2^{1/2}(\mathbb{T})}$  and  $|||\cdot|||_{W_2^{1/2}(\mathbb{T})}$  (see (2)), we infer that  $u_n \circ h \in W_2^{1/2}(\mathbb{T})$  for all  $n = 1, 2, \dots$ , and

$$\|u_n \circ h\|_{W_2^{1/2}(\mathbb{T})} \leq c \|u \circ h\|_{W_2^{1/2}(\mathbb{T})}, \quad n = 1, 2, \dots, \quad (10)$$



where  $c > 0$  does not depend on  $n$ .

The property of a function to be of bounded variation is invariant under homeomorphic changes of variable, hence from (7) it follows that  $u_n \circ h \in V(\mathbb{T})$  for all  $n$ . Certainly we also have  $u_n \circ h \in C(\mathbb{T})$ . Applying Lemma 4, and taking (10) into account, we obtain

$$\begin{aligned} \left| \frac{1}{2\pi} \int_{\mathbb{T}} v(t) du_n(t) \right| &= \left| \frac{1}{2\pi} \int_{\mathbb{T}} v \circ h(t) du_n \circ h(t) \right| \leq \\ &\leq \|v \circ h\|_{W_2^{1/2}(\mathbb{T})} \|u_n \circ h\|_{W_2^{1/2}(\mathbb{T})} \leq c \|v \circ h\|_{W_2^{1/2}(\mathbb{T})} \|u \circ h\|_{W_2^{1/2}(\mathbb{T})}, \end{aligned}$$

which contradicts (8). The theorem is proved.

*Remarks.* 1. For  $s > 0$  consider the Sobolev space  $W_2^s(\mathbb{T})$  i.e., the space of all (integrable) functions  $f$  with

$$\sum_{k \in \mathbb{Z}} |\widehat{f}(k)|^2 |k|^{2s} < \infty.$$

As the author of this paper showed in [7, Corollary 3], for each compact family  $K$  in  $C(\mathbb{T})$  (or equivalently for each class  $\text{Lip}_\omega(\mathbb{T})$ ) there exists a homeomorphism  $h$  of  $\mathbb{T}$  such that  $f \circ h \in \bigcap_{s < 1/2} W_2^s(\mathbb{T})$  for all  $f \in K$  (for all  $f \in \text{Lip}_\omega(\mathbb{T})$ ).

2. There exists a real-valued  $f \in C(\mathbb{T})$  such that  $f \circ h \notin \bigcup_{s > 1/2} W_2^s(\mathbb{T})$  for every homeomorphism  $h$  of  $\mathbb{T}$ . This is a simple consequence of the inclusion  $\bigcup_{s > 1/2} W_2^s \cap C(\mathbb{T}) \subseteq A(\mathbb{T})$ , where  $A(\mathbb{T})$  is the Wiener algebra of absolutely convergent Fourier series, and a well-known result of Olevskii, that provides a negative answer to Lusin's rearrangement problem: there exists a real-valued  $f \in C(\mathbb{T})$  such that  $f \circ h \notin A(\mathbb{T})$  for every homeomorphism  $h$  ([8], see also [9]).

3. The function  $f(t) = \sum_{k \geq 0} 2^{-k/2} e^{i2^k t}$  is in  $\text{Lip}_{1/2}(\mathbb{T})$ , (see, e.g., [1, Ch. XI, Sec. 6]), but it is obvious, that  $f \notin W_2^{1/2}(\mathbb{T})$ ; thus  $\text{Lip}_{1/2}(\mathbb{T}) \not\subseteq W_2^{1/2}(\mathbb{T})$ . The author does not know if the assertion of the theorem proved in this paper holds for  $\omega(\delta) = \delta^{1/2}$ . At the same time there is no change of variable which will bring the whole class  $\text{Lip}_{1/2}(\mathbb{T})$  into  $W_2^{1/2}(\mathbb{T})$ ; a proof will be presented in another paper.

## References

1. Bari N. K., *Trigonometric series*, Fizmatgiz, Moscow, 1961. English transl.: Vols. I, II, Pergamon Press, Oxford, and Macmillan, New York, 1964.
2. Bohr H., Über einen Satz von J. Pál, *Acta Sci. Math. (Szeged)* **7** (1935), 129–135.
3. Goffman G., Nishiura T., Waterman D., *Homeomorphisms in Analysis*, Mathematical Surveys and Monographs v. 54, Amer. Math. Soc., 1997.
4. Jurkat W., Waterman D., Conjugate functions and the Bohr–Pál theorem, *Complex Variables* **12** (1989), 67–70.
5. Kahane J.-P., Quatre leçons sur les homéomorphismes du cercle et les séries de Fourier, in: *Topics in Modern Harmonic Analysis*, Vol. II, Ist. Naz. Alta Mat. Francesco Severi, Roma, 1983, 955–990.
6. Kahane J.-P., Katznelson Y., Séries de Fourier des fonctions bornée, *Studies in Pure Math.*, in Memory of Paul Turán, Budapest, 1983, pp. 395–410. (Preprint, Orsay, 1978.)
7. Lebedev V. V., Change of variable and the rapidity of decrease of Fourier coefficients, *Mat. Sbornik* **181** (1990), 1099–1113 (Russian). English transl.: Math. USSR-Sb. **70** (1991), 541–555. English transl. corrected by the author is available at:  
<http://arxiv.org/abs/1508.06673>
8. Olevskii A. M., Change of variable and absolute convergence of Fourier series, *Dokl. Akad. Nauk SSSR* **256** (1981), 284–288 (Russian). English transl.: *Soviet Math. Dokl.* **23** (1981), 76–79.
9. Olevskii A. M., Modifications of functions and Fourier series, *Uspekhi Mat. Nauk* **40** (1985), 157–193 (Russian). English transl.: *Russian Math. Surveys* **40** (1985), 181–224.
10. Olevskii A. M., Modifications of functions and Fourier series, *Theory of Functions and Approximations*, Proc. Second Saratov Winter School (Saratov, Jan 24–Feb 5, 1984), Part 1, Saratov. Gos. Univ., Saratov, 1986, pp. 31–43 (Russian).

11. Pál J., Sur les transformations des fonctions qui font converger leurs séries de Fourier, *C. R. Acad. Sci Paris* **158** (1914), 101–103.
12. Saakjan A. A., Integral moduli of smoothness and the Fourier coefficients of the composition of functions, *Mat. Sbornik* **110** (1979), 597–608 (Russian). English transl.: *Mathematics of the USSR-Sbornik* **38** (1981), 549–561.

Moscow Institute of Electronics and Mathematics,  
National Research University Higher School of Economics  
34 Tallinskaya St.  
Moscow, 123458, Russia  
E-mail address: *lebedevhome@gmail.com*